US Preventative Services Task Force: Recommendation Statement. JAMA 2021;326(12):1186-1191

Table 1. Clinical Risk Assessment for Preeclampsia^a

Risk level	Risk factors	Recommendation
High ^b	 History of preeclampsia, especially when accompanied by an adverse outcome Multifetal gestation Chronic hypertension Pregestational type 1 or 2 diabetes Kidney disease Autoimmune disease (ie, systemic lupus erythematous, antiphospholipid syndrome) Combinations of multiple moderate-risk factors 	Recommend low-dose aspirin if the patient has ≥1 of these high-risk factors
Moderate ^c	 Nulliparity Obesity (ie, body mass index >30) Family history of preeclampsia (ie, mother or sister) Black persons (due to social, rather than biological, factors)^d Lower income^d Age 35 years or older Personal history factors (eg, low birth weight or small for gestational age, previous adverse pregnancy outcome, >10-year pregnancy interval) In vitro conception 	Recommend low-dose aspirin if the patient has ≥2 moderate-risk factors
		Consider low-dose aspirin if the patient has 1 of these moderate-risk factors ^d
_ow	Prior uncomplicated term delivery and absence of risk factors	Do not recommend low-dose aspirin

^a Includes only risk factors that can be obtained from the patient medical history.

some more consistently than others. A combination of multiple moderate-risk factors may place a pregnant person at higher risk for preeclampsia.

^b Includes single risk factors that are consistently associated with the greatest risk for preeclampsia. Preeclampsia incidence would likely be at least 8% in a population of pregnant individuals having 1 of these risk factors.

^c These factors are independently associated with moderate risk for preeclampsia,

^d These factors are associated with increased risk due to environmental, social, and historical inequities shaping health exposures, access to health care, and the unequal distribution of resources, not biological propensities.