



SAFE REDUCTION OF PRIMARY CESAREAN BIRTHS: SUPPORTING INTENDED VAGINAL BIRTHS

READINESS

Every Patient, Provider and Facility

- Build a provider and maternity unit culture that values, promotes, and supports spontaneous onset and progress of labor and vaginal birth and understands the risks for current and future pregnancies of cesarean birth without medical indication.
- Optimize patient and family engagement in education, informed consent, and shared decision making about normal healthy labor and birth throughout the maternity care cycle.
- Adopt provider education and training techniques that develop knowledge and skills on approaches which maximize the likelihood of vaginal birth, including assessment of labor, methods to promote labor progress, labor support, pain management (both pharmacologic and non-pharmacologic), and shared decision making.

RECOGNITION AND PREVENTION

Every patient

- Implement standardized admission criteria, triage management, education, and support for women presenting in spontaneous labor.
- Offer standardized techniques of pain management and comfort measures that promote labor progress and prevent dysfunctional labor.
- Use standardized methods in the assessment of the fetal heart rate status, including interpretation, documentation using NICHD terminology, and encourage methods that promote freedom of movement.
- Adopt protocols for timely identification of specific problems, such as herpes and breech presentation, for patients who can benefit from proactive intervention before labor to reduce the risk for cesarean birth.

PATIENT
SAFETY
BUNDLE

Safe Reduction of
Primary Cesarean Births



RESPONSE

To Every Labor Challenge

- Have available an in-house maternity care provider or alternative coverage which guarantees timely and effective responses to labor problems.
- Uphold standardized induction scheduling to ensure proper selection and preparation of women undergoing induction.
- Utilize standardized evidence-based labor algorithms, policies, and techniques, which allow for prompt recognition and treatment of dystocia.
- Adopt policies that outline standard responses to abnormal fetal heart rate patterns and uterine activity.
- Make available special expertise and techniques to lessen the need for abdominal delivery, such as breech version, instrumented delivery, and twin delivery protocols.

REPORTING/SYSTEMS LEARNING

Every birth facility

- Track and report labor and cesarean measures in sufficient detail to: 1) compare to similar institutions, 2) conduct case review and system analysis to drive care improvement, and 3) assess individual provider performance.
- Track appropriate metrics and balancing measures, which assess maternal and newborn outcomes resulting from changes in labor management strategies to ensure safety.

PATIENT SAFETY BUNDLE

Safe Reduction of Primary Cesarean Births